arXiv:1703.09101v1 [math.AP] 27 Mar 2017

ON THE L^2-CRITICAL NONLINEAR SCHRODINGER EQUATION WITH AN INHOMOGENEOUS DAMPING TERM.

MOHAMAD DARWICH

Abstract. We consider the L^2-critical nonlinear Schrodinger equation with an inhomogeneous damping term. We prove that there exists an initial data such that the corresponding solution is global in $H^1(R^d)$ and we give the minimal time of the blow up for some initial data.
Résumé. On considere l'equation de Schrodinger nonlinéaire L^2-critique avec un terme d'amortissement non homogène. On montre qu'ils existent des données initiales tels que la solution est globale dans $H^1(\mathbb{R}^d)$ et on donne le temps minimal d'explosion pour quelques donnes initiales.

1. Section francaise abrégée

On montre dans cette note qu'ils existent des données initiales tels que la solution est globale dans $H^1(\mathbb{R}^d)$ et on donne le temps minimal d'explosion pour quelques donnes initiales, le resultat sera obtenue en montrant qu'un phénomène de L^2 concentration aura lieu proche de point d'explosion.
Notre résultat principal concernant l'equation (2.1)est donné dans le théorème suivant:

Theorem 1.1. *Soit $u_0 \in H^1(\mathbb{R}^d)$ et $d = 1, 2, 3, 4$:*

(1) *Si $a(x) > 0$ et $\|u_0\|_{L^2} \leq \|Q\|_{L^2}$, alors la solution de (2.1) est globale dans $H^1(\mathbb{R}^d)$.*

(2) *Si $a(x)$ est de signe quelconque et s'il existent des données initiales u_0 avec $\|u_0\|_{L^2} \leq \|Q\|_{L^2}$ tels que la solution de (2.1) explose en temps fini T alors $T > \frac{1}{\|a\|_{L^\infty}} \log(\frac{\|Q\|_{L^2}}{\|u_0\|_{L^2}})$.*

Key words and phrases. Damped Nonlinear Schrödinger Equation, Blow-up, Global existence.

2. Introduction

In this paper, we study the Cauchy problem for the L^2-critical nonlinear Schrödinger equations:

$$\begin{cases} iu_t + \Delta u + |u|^{\frac{4}{d}}u + ia(x)u = 0, (t,x) \in [0,\infty[\times\mathbb{R}^d, d = 1,2,3,4. \\ u(0) = u_0 \in H^1(\mathbb{R}^d) \end{cases} \tag{2.1}$$

with a real inhomogeneous damping term a $\in C^1(\mathbb{R}^d,\mathbb{R}) \cap W^{1,\infty}(\mathbb{R}^d,\mathbb{R})$ and initial data $u(0) = u_0 \in H^1(\mathbb{R}^d)$.Equation (2.1) arises in several areas of nonlinear optics and plasma physics. The inhomogenous damping term corresponds to an electromagnetic wave absorved by an inhomogenous medium. (cf [1], [2])
It is known that the Cauchy problem for (2.1) is locally well-posed in $H^1(\mathbb{R}^d)$(see Kato [8] and also Cazenave [4]): For any $u_0 \in H^1(\mathbb{R}^d)$, there exist $T \in (0,\infty]$ and a unique solution $u(t)$ of (2.1) with $u(0) = u_0$ such that $u \in C([0,T); H^1(\mathbb{R}^d))$. Moreover, T is the maximal existence time of the solution $u(t)$ in the sense that if $T < \infty$ then $\lim\limits_{t\to T} \|u(t)\|_{H^1(\mathbb{R}^d)} = \infty$.
Dias and figueira [6] studied the supercritical case($|u|^p u$ with $p > \frac{4}{d}$) and showed that blow-up in finite time can occur, using the virial method. In [5], Correia was studied the equation in dimension one, and he proved the existence of blowup phenomena in the energy space H^1
Let us notice that for $a = 0$ (2.1) becomes the L^2-critical nonlinear Schrödinger equation:

$$\begin{cases} iu_t + \Delta u + |u|^{\frac{4}{d}}u = 0 \\ u(0) = u_0 \in H^1(\mathbb{R}^d) \end{cases} \tag{2.2}$$

Special solutions play a fundamental role for the description of the dynamics of (2.2). They are the solitary waves of the form $u(t,x) = \exp(it)Q(x)$, where Q solves:

$$\Delta Q + Q|Q|^{\frac{4}{d}} = Q. \tag{2.3}$$

Let u be a solution of (2.1), we define the following quantities:
L^2norm : $\|u(t)\|_{L^2}$.
Energy : $E(u(t)) = \frac{1}{2}\|\nabla u\|_{L^2}^2 - \frac{d}{4+2d}\|u\|_{L^{\frac{4}{d}+2}}^{\frac{4}{d}+2}$.
Kinetic momentum : $P(u(t)) = Im(\int \nabla u\overline{u}(t,x))$.

It is easy to prove that if u is a solution of (2.1) then :

$$\frac{d}{dt}\|u(t)\|_{L^2}^2 = -\int a(x)|u(t,x)|^2 dx, t \in [0,T), \tag{2.4}$$

$$\frac{d}{dt}E(u(t)) = -\int a(x)\nabla u(t,x)dx + \int a(x)u(t,x)^{\frac{4}{d}+2} - \Re\int(\nabla u.\nabla a)\overline{u}dx \tag{2.5}$$

and

$$\frac{d}{dt}P(u(t)) = -2\int a(x)\Im(\nabla u\overline{u})dx, t\in[0,T). \tag{2.6}$$

Remark 2.1. *Remark that if $a(x) > 0, \forall x \in \mathbb{R}^d$, then $\|u(t)\|_{L^2} < \|u_0\|_{L^2}$, for all $t \in [0,T[$.*

Let us now our results:

Theorem 2.1. *Let $u_0 \in H^1(\mathbb{R}^d)$ with $d = 1,2,3,4$, then:*

1. *If $a(x) > 0$ and $\|u_0\|_{L^2} \leq \|Q\|_{L^2}$, then the corresponding solution of (2.1) is global in H^1.*
2. *If $a(x)$ has an arbitrary sign and if there exists an initial data $u_0 \in H^1$ with $\|u_0\| < \|Q\|_{L^2}$ such that the corresponding solution blows up at finite time T, then $T > \frac{1}{\|a\|_\infty}\log(\frac{\|Q\|_{L^2}}{\|u_0\|_{L^2}})$.*

3. L^2-CONCENTRATION

In this section, we prove theorem 2.1 by extending the proof of the L^2-concentration phenomen, proved by Ohta and Todorova [10] in the radial case, to the non radial case.
Hmidi and Keraani showed in [7] the L^2-concentration for the equation (2.2) without the hypothese of radiality, using the following theorem:

Theorem 3.1. *Let $(v_n)_n$ be a bounded family of $H^1(\mathbb{R}^d)$, such that:*

$$\limsup_{n\to+\infty}\|\nabla v_n\|_{L^2(\mathbb{R}^d)} \leq M \quad \text{and} \quad \limsup_{n\to+\infty}\|v_n\|_{L^{\frac{4}{d}+2}} \geq m. \tag{3.1}$$

Then, there exists $(x_n)_n \subset \mathbb{R}^d$ such that:

$$v_n(\cdot + x_n) \rightharpoonup V \quad \text{weakly},$$

with $\|V\|_{L^2(\mathbb{R}^d)} \geq (\frac{d}{d+4})^{\frac{d}{4}}\frac{m^{\frac{d}{2}+1}+1}{M^{\frac{d}{2}}}\|Q\|_{L^2(\mathbb{R}^d)}$.

Now we have the following theorem:

Theorem 3.2. *Assume that $u_0 \in H^1(\mathbb{R}^d)$, and suppose that the solution of (2.1) with $u(0) = u_0$ blows up in finite time $T \in (0,+\infty)$. Then, for any function $w(t)$ satisfying $w(t)\|\nabla u(t)\|_{L^2(\mathbb{R}^d)} \to \infty$ as $t \to T$, there exists $x(t) \in \mathbb{R}^d$ such that, up to a subsequence,*

$$\limsup_{t\to T}\|u(t)\|_{L^2(|x-x(t)|<w(t))} \geq \|Q\|_{L^2(\mathbb{R}^d)}.$$

To show this theorem we shall need the following lemma:

Lemma 3.1. *Let $T \in (0, +\infty)$, and assume that a function $F : [0, T) \longmapsto (0, +\infty)$ is continuous, and $\lim_{t\to T} F(t) = +\infty$. Then, there exists a sequence $(t_k)_k$ such that $t_k \to T$ and*

$$\lim_{t_k\to T} \frac{\int_0^{t_k} F(\tau)d\tau}{F(t_k)} = 0. \tag{3.2}$$

For the proof see [10].

Proof of Theorem 3.2:
Suppose that there exist an initial data u_0 in H^1 with $\|u_0\| \leq \|Q\|_{L^2}$ such that the corresponding solution blows up at finite time T.
By the energy identity (2.5), we have

$$E(u(t)) = E(u_0) - \int_0^t H(u(\tau))d\tau, \quad t \in [0, T[. \tag{3.3}$$

Where $H(u(t)) = -\int a(x)|\nabla u(t,x)|^2dx + \int a(x)|u(t,x)|^{\frac{4}{d}+2} - \Re\int(\nabla u.\nabla a)\overline{u}dx$.
Let us recall the Gagliardo-Nirenberg inequality:

$$\|u\|^{2+\frac{4}{d}}_{L^{2+\frac{4}{d}}} \leq C\|\nabla u\|^2_{L^2}\|u\|^{\frac{4}{d}}_{L^2}, \tag{3.4}$$

Note that (2.4) gives that:

$$\|u_0\|_{L^2}e^{-\|a\|_{L^\infty}t} \leq \|u\|_{L^2} \leq \|u_0\|_{L^2}e^{\|a\|_{L^\infty}t} \tag{3.5}$$

Now using (3.4) and (3.5) we obtain that:

$$\begin{aligned}|H(u(t))| &\leq \|a\|_{L^\infty}\|\nabla u(t)\|^2_{L^2(\mathbb{R}^d)} + \|a\|_{L^\infty}\|u(t)\|^{2+\frac{4}{d}}_{L^{2+\frac{4}{d}}} + \|\nabla a\|_{L^\infty}\|u\|_{L^2}\|\nabla u\|_{L^2}\\ &\leq \|a\|_{L^\infty}\|\nabla u(t)\|^2_{L^2(\mathbb{R}^d)} + C\|a\|_{L^\infty}e^{\|a\|_{L^\infty}t}\|\nabla u\|^2_{L^2}\|u_0\|^{\frac{4}{d}}_{L^2}\\ &+ e^{\|a\|_{L^\infty}t}\|\nabla a\|_{L^\infty}\|u_0\|_{L^2}\|\nabla u\|_{L^2}\end{aligned}$$

for all $t \in [0, T[$. Then

$$|H(u(t))| \leq C(\|a\|_{H^1}, \|u_0\|_{L^2})e^{\|a\|_{L^\infty}t}\|\nabla u\|^2_{L^2}$$

Moreover, we have $\lim_{t\to T}\|\nabla u(t)\|_{L^2(\mathbb{R}^d)} = +\infty$, thus by Lemma 3.1, there exists a sequence $(t_k)_k$ such that $t_k \to T$ and

$$\lim_{k\to\infty} \frac{\int_0^{t_k} H(u(\tau))d\tau}{\|\nabla u(t_k)\|^2_{L^2(\mathbb{R}^d)}} = 0. \tag{3.6}$$

Let

$$\rho(t) = \frac{\|\nabla Q\|_{L^2(\mathbb{R}^d)}}{\|\nabla u(t)\|_{L^2(\mathbb{R}^d)}} \quad \text{and} \quad v(t,x) = \rho^{\frac{d}{2}}u(t,\rho x)$$

and $\rho_k = \rho(t_k), v_k = v(t_k, .)$. The family $(v_k)_k$ satisfies

$$\|v_k\|_{L^2(\mathbb{R}^d)} \leq e^{\|a\|_{L^\infty} T} \|u_0\|_{L^2(\mathbb{R}^d)} \quad \text{and} \quad \|\nabla v_k\|_{L^2(\mathbb{R}^d)} = \|\nabla Q\|_{L^2(\mathbb{R}^d)}.$$

By (3.3) and (3.6), we have

$$E(v_k) = \rho_k^2 E(u_0) - \rho_k^2 \int_0^{t_k} H(u(\tau))d\tau \to 0, \tag{3.7}$$

which yields

$$\|v_k\|^{\frac{4}{d}+2}_{L^{\frac{4}{d}+2}} \to \frac{d+2}{d} \|\nabla Q\|^2_{L^2(\mathbb{R}^d)}. \tag{3.8}$$

The family $(v_k)_k$ satisfies the hypotheses of Theorem 3.1 with

$$m^{\frac{4}{d}+2} = \frac{d+2}{d} \|\nabla Q\|^2_{L^2(\mathbb{R}^d)} \quad \text{and} \quad M = \|\nabla Q\|_{L^2(\mathbb{R}^d)},$$

thus there exists a family $(x_k)_k \subset \mathbb{R}^d$ and a profile $V \in H^1(\mathbb{R}^d)$ with $\|V\|_{L^2(\mathbb{R}^d)} \geq \|Q\|_{L^2(\mathbb{R}^d)}$, such that,

$$\rho_k^{\frac{d}{2}} u(t_k, \rho_k \cdot + x_k) \rightharpoonup V \in H^1 \quad \text{weakly}. \tag{3.9}$$

Using (3.9), $\forall A \geq 0$

$$\liminf_{n\to+\infty} \int_{B(0,A)} \rho_n^d |u(t_n, \rho_n x + x_n)|^2 dx \geq \int_{B(0,A)} |V|^2 dx,$$

but $\lim_{n\to+\infty} \frac{w(t_n)}{\rho_n} = +\infty$ thus $\frac{w(t_n)}{\rho_n} > A$, $\rho_n A < w(t_n)$. This gives immediately:

$$\liminf_{n\to+\infty} \sup_{y\in\mathbb{R}^d} \int_{|x-y|\leq w(t_n)} |u(t_n, x)|^2 dx \geq \int_{|x|\leq A} |V|^2 dx.$$

This it is true for all $A > 0$ thus :

$$\liminf_{t\to T} \sup_{y\in\mathbb{R}^d} \int_{|x-y|\leq w(t)} |u(t, x)|^2 dx \geq \int Q^2 dx. \tag{3.10}$$

Finally we obtain:

(1) If $a(x) > 0$, then the norm L^2 is strictly decreasing, with (3.10) in hand we obtain that:$\|u_0\|^2_{L^2} > \|Q\|^2_{L^2}$

(2) If the sign of a is arbitrary, (3.10) gives that $e^{2\|a\|_{L^\infty} T} \|u_0\|^2_{L^2} \geq \|Q\|^2_{L^2}$.

Proof of Theorem 2.1:
Now if $a(x) > 0$ $\forall x \in \mathbb{R}^d$, and u_0 be an initial data such that $\|u_0\| \leq \|Q\|$, we obtain a contradiction, that means that the solution is global in H^1, and this gives the proof of part 1 of the theorem.

If a has an arbitrary sign, and if u blows up with initial data u_0 with $\|u_0\|_{L^2} < \|Q\|_{L^2}$ at finite time T, we obtain that $T > \frac{1}{\|a\|_{L^\infty}} \log(\frac{\|Q\|_{L^2}}{\|u_0\|_{L^2}})$, this gives the proof of the second part of the theorem.

References

[1] G. Barontini, R. Labouvie, F. Stubenrauch, A. Vogler, V. Guarrera and H. Ott, Controlling the dynamics of an open many-body quantum system with localized dissipation, Phys. Rev. Lett., 110 (2013), 35302-35305.

[2] V.A. Brazhnyi, V.V. Konotop, V.M. Prez-Garcia and H. Ott, Dissipation-Induced coherent structures in Bose-Einstein Condensates, Phys. Rev. Lett., 102 (2009), 144101-144104.

[3] H. Berestycki and P.-L. Lions. *Nonlinear scalar field equations. II. Existence of infinitely many solutions*. Arch. Rational Mech. Anal., 82(1983):347–375.

[4] T. Cazenave. *Semilinear Schrödinger equations*, volume 10 of *Courant Lecture Notes in Mathematics*. New York University Courant Institute of Mathematical Sciences, New York, 2003.

[5] S.Correia *Blowup for the nonlinear Schrodinger equation with an inhomogeneous damping term in the L^2-critical case*.https://arxiv.org/pdf/1410.8011.pdf

[6] J.P Dias and M. Figueira. *On the blowup of solutions of a Shrodinger equation with an inhomogeneous damping coefficient* Communications in Contemporary Mathematics Volume 16, Issue 03, June 2014.

[7] T. Hmidi and S. Keraani. *Blowup theory for the critical nonlinear Schrödinger equations revisited*. Int. Math. Res. Not., 46(2005):2815–2828.

[8] T. Kato. *On nonlinear Schrödinger equations* Ann. Inst. H. Poincaré Phys. Théor., 46(1987):113–129.

[9] M.K Kwong. *Uniqueness of positive solutions of $\Delta u - u + u^p = 0$ in $\mathbf{R}^n$*. Arch. Rational Mech. Anal., 105(1989):243–266.

[10] M. Ohta and G. Todorova. *Remarks on global existence and blowup for damped nonlinear Schrödinger equations*. Discrete Contin. Dyn. Syst., 23(2009):1313–1325.

[11] M. Tsutsumi. *Nonexistence of global solutions to the Cauchy problem for the damped nonlinear Schrödinger equations*. SIAM J. Math. Anal., 15(1984):357–366.

[12] M.I. Weinstein. *Nonlinear Schrödinger equations and sharp interpolation estimates*. Comm. Math. Phys., 87(1982/83):567–576.

M. Darwich: Lebanese University, Hadath.
E-mail address: Mohamad.Darwich@lmpt.univ-tours.fr